Cucumber Beetle Lure Manual

7030 E 46th Avenue Dr, Denver, CO 80216, United States
support@hensleyseeds.com | hensleyseeds.com

Cucumber beetles are a major problem wherever cucurbits, such as squash, cucumbers, watermelon, and pumpkins, are grown. They can kill or severely stunt seedlings; damage stems and fruits on older plants; and roots can be damaged by beetle larvae. It is well known for transmitting a range of diseases.

Place one lure inside a Sticky Yellow Trap and hang upwind of the crop to be monitored. Use one trap with lure per 2,000 square feet for monitoring or per 400 square feet for trapping. Replace when filled with beetles or dirty. The lure is effective for 3–4 weeks, depending on temperature — the attractant dissipates faster in warmer conditions, reducing the length of effective control.



Mature cucumber beetle

REV 10/11/2015 RC, MN

Copyright © 2016 Hensley's Seed Co. All rights reserved.

Cucumber Beetle Lure Manual

7030 E 46th Avenue Dr, Denver, CO 80216, United States
support@hensleyseeds.com | hensleyseeds.com

Cucumber beetles are a major problem wherever cucurbits, such as squash, cucumbers, watermelon, and pumpkins, are grown. They can kill or severely stunt seedlings; damage stems and fruits on older plants; and roots can be damaged by beetle larvae. It is well known for transmitting a range of diseases.

Place one lure inside a Sticky Yellow Trap and hang upwind of the crop to be monitored. Use one trap with lure per 2,000 square feet for monitoring or per 400 square feet for trapping. Replace when filled with beetles or dirty. The lure is effective for 3–4 weeks, depending on temperature — the attractant dissipates faster in warmer conditions, reducing the length of effective control.



Mature cucumber beetle

REV 10/11/2015 RC, MN

Copyright © 2016 Hensley's Seed Co. All rights reserved.

1

HENSLEY'S SEED CO. | GROWING KNOWLEDGE FOR EVERY GARDENER